# Adverse skin reactions of personal protective equipment on health care workers against COVID -19

Santosh M. Soraganvi Blood Bank Assistant

Lions Blood bank Vijayapur, Karnataka.

Revanshiddayya S Hiremath Laboratory technologists,

Belagavi Institute of Medical Sciences, Belagavi.

Panchaksharayya S Hiremath Laboratory technologists,

Belagavi Institute of Medical Sciences, Belagavi.

Manjunath D. Marad PhD scholar

Dept of Community Medicine Shri B M Patil Medical

College, Hospital & DE (deemed to

be University) Vijayapur.

Eknath P Jadhav Assistant Hospital Administrator,

Shri B M Patil Medical College, Hospital & Dr; Research

Centre BLDE (deemed to be University) Vijayapur.

Gavishiddappa A Hadimani Assistant Professor,

Dept of Anatomy,

Shri B M Patil Medical College, Hospital & Dr; Research

Centre BLDE (deemed to be University) Vijayapur.

The coronavirus outbreak, which was first detected in China, has infected people in 185 countries. COVID-19 is a pneumonia-like disease caused by a new coronavirus, named SARS-CoV-2, that is similar to the virus that causes Severe Acute Respiratory Syndrome (SARS Coronavirus disease 2019 (COVID-19). To study the prevalence of adverse reactions of wearing PPE among healthcare workers in Karnataka India during the Covid -19 outbreak. Twenty-one-day observation were made on twenty laboratory health workers and observed hazardous skin effect on health workers wearing PPE long hours during working period. long hour wearing of PPE in our study found respiratory complications, sweating and skin damage. Adequate dermatological knowledge of PPE use and design and extensive comprehensive training is important in risk of pandemic outbreak.

Keywords: Corona Virus, Covid -19, Personal Protective Equipment, Transmission

#### Introduction

Coronaviruses are important human and animal pathogens. Corona viruses are a large family of enveloped, non-segmental, positive sensed single stranded RNA Viruses. Measures around 120 diameter [1]. Novel corona virus (SARSCOV-2) is currently causing a severe outbreak of disease (termed COVID -19) in China and multiple other countries, threatening to cause global pandemic [2,3]. In humans, single stranded RNA CoVs mostly causes respiratory and gastrointestinal symptoms. Clinical manifestations range from a mild flu-like symptoms to more severe disease such as bronchitis pneumonia, severe acute respiratory distress syndrome, multi organ failure and even leads to death [4]. This is the third extremely pathogenic human coronavirus which has emerged recently after severe acute respiratory syndrome (SARS). The World Health Organisation (WHO)

1/5

has declared the COVID-19 outbreak as a global pandemic. Research Studies on corona have suggested that COVID-19 can have high transmission rates with the average infected person able to spread the disease to two or three other individuals [5]. Worldwide, as millions of people stay at home to minimise transmission of severe acute respiratory syndrome coronavirus -2, Health care workers are at an especially high risk of catching the <u>coronavirus</u>, because of their prolonged exposure to confirmed Covid -19 patients. Personal protective equipment plays crucial role in prevention of transmission of infection in the health care setting

#### **Aims & Objectives**

To study the prevalence of adverse reactions of wearing PPE among healthcare workers in Karnataka India during the Covid -19 outbreak.

# **Material and Methods**

Study was carried out on twenty health workers of civil hospital Belgaum Karnataka state India. Study was started from 15<sup>th</sup> March 2020 to 6<sup>th</sup> May 2020. Observations were made up to twenty-one days. Before observational study all the health workers were underwent for comprehensive training program and learnt how to don (put on) and doff (take off) PPE, limitations of PPE, proper care, maintenance, and disposal of PPE. PPE kit includes gloves, medical masks, goggles or a face shield, and gowns. In our study we observed hazardous skin effect on health workers wearing PPE long hours during working period.

#### **Results**

Personal protective equipment (PPE) has been essential to protecting healthcare workers from the novel coronavirus, but it presents its own set of challenges to their health, notably respiratory complications, sweating and skin damage associated with prolonged use of PPE. During our study we found significant effects on health workers wearing long period PPE kit they were

- 1. Wearing mask problems: Out of twenty, fourteen health worker developed skin damage by wearing mask for long hours in laboratory. Enhanced use of mask can cause redness maceration erythema papule scaling and even colour changes in prolonged use cases particularly on the bridge of the nose and top of the cheeks, and the belt of the mask causes pain on the auricular cartilage. Most of the health worker in our study complained breathlessness, burning, stinging, itching acne and even dermatitis over the face (figure1.g).
- 2. Wearing gloves: Most of the laboratory health worker complained irritant contact dermatitis, allergic contact dermatitis, contact urticaria hyper hydration of skin, maceration erosion and eczema type of symptoms. (figure1.a, c, d & h).
- 3. Face shield: Significantly less number of (6/20) health worker developed skin rash acne and red ness sweating over the forehead region due to compression of belt on forehead for long period in laboratory. In most of health worker we observed glare fogging and impaired vision during working in laboratory.
- 4. Wearing gown: Most of the health worker developed heavy sweating, skin rashes, acnes particularly on the front and back of the chest, pain, Itching, prickling, and fungal infection after wearing PPE gown for long hour. (figure 1.b, e f &, i).

Table 1 Showing the skin problem of Health care workers, working in the civil hospital Covid-19 laboratory

Sr no	PPE Kit	Health worker (n=20)	Skin effect

1	Mask		Ear Pain Sweating skin rash, breathlessness
2	Goggle		Sweating, nose pad and temple markings
3	Gloves	•	Skin rashes. Maceration, and erosion
4	Face shield		Sweating, Impaired vision, glare, fogging
5	Protective Gowns	18/20	Sweating, heat, skin infection

Table 1. The skin problem of Health care workers, working in the civil hospital Covid-19 laboratory

Figure 1 Showing the skin problem of Health care workers, working in the civil hospital Covid-19 laboratory




Figure 1. The skin problem of Health care workers, working in the civil hospital Covid-19 laboratory

#### **Discussion**

For COVID -19 exposure, PPE is specialized protective equipment used to prevent contact with hazardous substances. Its use is an integral part of infection control and prevention measures that protect workers from exposure to blood, body fluids, and other potentially infectious materials [6].

PPE such as gowns, gloves, masks, and goggles provide physical barriers that prevent the hands, skin, clothing, eyes, nose, and mouth from coming in contact with infectious agents. Plays prominent role to ensuring overall health and safety of health workers, by reducing exposure to transmission of COVID-19[7].

Masks Plays an essential role in the fight against the COVID-19 pandemic especially for health workers. Masks should be good quality and designed with advanced technology to seal around the

nose and mouth to achieve very close facial fit and very efficient filtration of air born particles to reduce the chance of breathlessness and other skin infection [8].

Gloves are very important component personal protective equipment and act as a protective barrier to prevent the possible transmission of Covid-19 for health care workers, sweat can make wearing gloves uncomfortable and even lead to skin problems that make the issue worse. When skin is exposed to sweat for a prolonged time, it weakens and becomes more vulnerable; the moist environment is the perfect breeding ground for bacteria and fungus [9]. To avoid such type of complication health worker should change the gloves after every one hour may prevent the skin infection .

Face shields are important protective barrier for the facial area and associated mucous membranes from airborne body fluids (blood, saliva, bronchial secretions, vomit etc.) expelled as a result of various physiological processes (vomiting, coughing, sneezing, etc.) and medical, dental, and veterinary procedures (suctioning the airway, placing nasogastric tubes, obstetrical procedures, surgery, dental procedures, etc.) [10]. Face shields should be efficacious and must fit snugly to afford a good seal to the forehead and should cover complete area of face and chin to prevent slippage of the device. Face shields should be selected that have visors treated for anti-glare, anti-static, and anti-fogging properties.

PPE gown is most important barrier to prevent transmission of pandemic outbreak of COVID -19 by blocking the transmission of contaminants from blood, body fluids, or respiratory secretions and it is essential for frontline health care workers [11]. PPE should be of good quality and waterproof to reduce the chance of skin infection. Effective use of PPE includes properly removing and disposing of contaminated PPE to prevent exposing both the wearer and other people to infection.

## **Conclusion**

The use of personal protective equipment (PPE) is an important strategy to protect healthcare personnel from contamination and to prevent the spread of pathogens to subsequent patients. Adequate dermatological knowledge of PPE use and design and extensive comprehensive training is important in risk of pandemic outbreak.

### References

- 1. WHO Emergencies preparedness, response, Novel Coronavirus China Disease outbreak news: Update 12 January 2020 https://www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en/.
- 2. Fehr AR, Perlman S. Coronaviruses: an overview of their replication and pathogenesis. Methods Mol Biol. 2015; 1282:1-23. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4369385/
- 3. Denison MR, Graham RL, Donaldson EF, Eckerle LD, Baric RS. Coronaviruses: an RNA proofreading machine regulates replication fidelity and diversity. RNA Biol. 2011;82::270-9. https://pubmed.ncbi.nlm.nih.gov/21593585/
- 4. Bagoji I B, & Bharatha A. COVID -19 and Robotics-creativities spark in the adverse. South East Asia Journal of Medical Sciences, 2020;41: 1. https://littlebaypublishers.com/index.php/seajournalms/article/view/61/53. file:///C:/Users/Ambadasu/Downloads/61-246-2-PB%20(3pdf
- 5. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. journal of advance research 2020; 24: 91-98. https://www.sciencedirect.com/science/article/pii/S2090123220300540
- 6. Verbeek JH, Rajamaki B, Ijaz S, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. Cochrane Database Syst Rev. 2020;4:CD011621.

- https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD011621.pub4/full
- 7. Roberto L G. The Severe Acute Respiratory Syndrome Coronavirus-2 (SARS CoV-2) in Dentistry. Management of Biological Risk in Dental Practice. Int. J. Environ. Res. Public Health 2020, 17: 3067. file:///C:/Users/Ambadasu/Downloads/ijerph-17-03067%20(1).pdf
- 8. Adams JG, Walls RM. Supporting the Health Care Workforce During the COVID-19 Global Epidemic. JAMA. 2020;32315:1439-1440. https://europepmc.org/article/med/32163102
- 9. Drascovics SW. Single-Use Gloves: Problems and Solutions. https://www.chemengonline.com/single-use-gloves-problems-solutions/
- 10. Roberge RJ. Face shields for infection control: A review. J Occup Environ Hyg. 2016;134:235-42. https://pubmed.ncbi.nlm.nih.gov/26558413/
- 11. World Health Organization. Rational use of personal protective equipment (PPE) for coronavirus disease (COVID-19): interim guidance. https://apps.who.int/iris/handle/10665/331498 (Updated on March 19, 2020).

5/5